

**Fig. 1** Map of the Sanya Bay coast showing the sampling locations ( ).

concentration of 1%.

Species were observed under an inverted microscope (OLYMPUS, IX71), and if necessary, Calcofluor White M2R at a final concentration of 10  $\mu\text{g/ml}$  was used to stain the outer shells of the dinoflagellates<sup>14</sup>, using an inverted differential interference contrast epifluorescence microscope (OLYMPUS, BX60F) equipped with a UV excitation unit. Photographs of these algae were captured with a charge-coupled device camera (OLYMPUS, DP70). *Prorocentrum* spp. were identified according to previous literatures<sup>3, 12, 15, 16</sup>.

## RESULTS

The genus *Prorocentrum* Ehrenberg, which is a desmokont dinoflagellate, characterized by apical insertion of the flagella, absence of the cingulum and the sulcus<sup>9</sup>, and the cell consists of the two lateral thecae, which are joined by the marginal edges<sup>12</sup>. In the present study, five species were from the coastal areas along Sanya Bay as described below.

### *P. hoffmannianum* Faust 1990

Synonym: *Exuviaella hoffmannianum* (Faust) McLachlan et Boalch 1997

Cell is ovoid, broadest in the middle region, tapering slightly apically, 45–55  $\mu\text{m}$  in length and 40–45  $\mu\text{m}$  in width, containing golden-brown chloroplasts, a centrally located pyrenoid, and a posterior nucleus. The valves are apically excavated. The periflagellar area is a wide triangle situated apically on the right valve, and it lacks both valve spines and anterior spines (Fig. 2a).

*P. hoffmannianum* is a benthic species, can be tycho planktonic, with warm temperate to tropical coastal waters distribution, reported from Belize<sup>17</sup>, and this species was reported to be associated with floating detritus in tropical coastal regions of the Caribbean Sea<sup>17</sup>, attached to macroalgae in the Belizean barrier reef ecosystem<sup>18</sup>.

*P. hoffmannianum* is usually considered toxic, producing fast-acting toxin and okadaic acid<sup>19</sup>.

### *P. lima* (Ehrenberg) Stein 1878

Basionym: *Cryptomonas lima* Ehrenberg 1860

Synonyms: *E. marina* Cienkowski 1881

*Dinopyxis laevis* Stein 1883

*E. lima* (Ehrenberg) Bütschli 1885

*E. lima* (Ehrenberg) Schütt 1896

*E. laevis* (Stein) Schroder 1900

*E. cincta* Schiller 1918

*E. marina* var. *lima* (Ehrenberg) Schiller 1931

*E. ostenfeldi* Schiller 1933

*E. caspica* Kiselev 1940

*P. marinum* Dodge et Bibby 1973

*P. arenarium* Faust 1994

Cells are oblong to ovate, broad in the middle region, narrow at the anterior end, 31–47  $\mu\text{m}$  in length and 22–40  $\mu\text{m}$  in width (Fig. 2b, 2c), containing a central pyrenoid and a large posterior nucleus (Fig. 2b). The valve surface is covered with scattered pores, but the centre is devoid of the pores, the right valve having valve pores of two different sizes, and the marginal pores are presented (Fig. 2c). The periflagellar area is a wide triangle containing a curved apical collar, void of valve spines or anterior spines (Fig. 2b).


*P. lima* is a neritic and estuarine, benthic/epiphytic species (can be tycho planktonic), worldwide distribution, typically in temperate and tropical waters<sup>8, 20</sup>, reported from East China Sea and South China Sea<sup>21</sup>; Japan, Indonesia and Philippines<sup>22</sup>, and this species was reported to be attached to macroalgae, or was observed swimming close to the bottom substrate, and was associated with coral reefs<sup>17</sup>, or can be found attached to floating detritus in mangrove habitats<sup>23</sup>.

*P. lima* is a toxic species known to produce a number of toxic substances, such as fast-acting toxin<sup>24</sup>, and diarrhoeic shellfish poisoning (DSP) toxins including okadaic acid<sup>25</sup>.

### *P. micans* Ehrenberg 1834

Synonyms: *P. schilleri* Böhrn 1933

*P. levantinoides* Bursa 1959



**Fig. 2** Light and fluorescence microphotographs of *Prorocentrum* spp.; (a) *P. hoffmannianum*, (b, c) *P. lima*, (d, e) *P. micans*, (f, g) *P. rhathymum*, (h, i) *P. sigmoides*. Scale bars 10  $\mu\text{m}$ .

#### *P. pacificum* Wood 1963

Cells are tear-drop to heart shaped, generally rounded anteriorly, tapering posteriorly, broadest around the middle, 35–70  $\mu\text{m}$  in length and 20–50  $\mu\text{m}$  in width (Fig. 2d, 2e). One convex side and one arched side, the convex arch profile is typically in the middle of the cell where is the broadest, and a well-developed winged apical spine is presented (Fig. 2d, 2e). Numerous tubular trichocyst pores are presented in short rows arranged radially (Fig. 2d). The periflagellar area is a relatively small, shallow, broad triangular depression situated apically on the right valve off-centre (Fig. 2e).

*P. micans* is a neritic and estuarine, planktonic species, but also found in oceanic environments, with cold temperate to tropical waters distribution<sup>20</sup>, reported from Japan<sup>16</sup>; Mexican Pacific Coast<sup>13</sup>; East China Sea and South China Sea<sup>21</sup>.

Although *P. micans* is capable of forming extensive blooms, it is usually considered harmless<sup>26</sup>. It may excrete substances that inhibit diatom growth, but apparently these substances do not enter the food chain or affect organisms at higher trophic levels<sup>27</sup>. Early reports on *P. micans* being a paralytic shellfish poison producer<sup>28</sup> are unconfirmed, and recent incidents involving shellfish mortality have

been attributed to oxygen depletion<sup>29</sup>.

#### *P. rhathymum* Loeblich III, Sherley et Schmidt 1979

Cells are asymmetric, ovate to oblong with straight sides, 30–38  $\mu\text{m}$  in length and 20–25  $\mu\text{m}$  in width (Fig. 2f, 2g). The valve pores form the line patterns in the posterior part of the valve (Fig. 2f), the marginal pores are presented (Fig. 2g), and the centres of the valves are devoid of pores (Fig. 2f, 2g). The periflagellar area, located apically and off-centre on the right valve, is a relatively small, V-shaped, shallow depression, and the curved periflagellar collar may appear as an apical spine (Fig. 2f, 2g).

*P. rhathymum* is a neritic and estuarine, benthic species, can be tycho planktonic, with tropical and subtropical waters distribution<sup>20</sup>, reported from New Caledonia and Ryukyu Islands (Japan)<sup>30</sup>, and this species was reported to be associated with microalgae, dead coral and seagrasses in Malaysia<sup>31</sup>.

*P. rhathymum* may produce toxins with haemolytic activity<sup>32</sup>, and water-soluble fast-acting toxin<sup>24</sup>.

### *P. sigmoides* Böhm 1933

Cells are elongated, in 'S' shape, one convex side and one generally straight side, and tapering posteriorly, 60–85  $\mu\text{m}$  in length and 20–30  $\mu\text{m}$  in width (Fig. 2h, 2i). Numerous trichocyst pores are presented (Fig. 2h, 2i). A sharp and tiny apical spine protruded from the anterior side, adjacent to the periflagellar area, sometimes slightly sigmoid, with the presence of an antapical mucron (Fig. 2i).

*P. sigmoides* is a planktonic species, with temperate to tropical waters distribution<sup>16</sup>, reported from South China Sea and Japan<sup>21</sup>.

*P. sigmoides* has never been reported to be a toxin producer, but it is a fish killer<sup>33</sup>, capable of forming extensive blooms, which can consume dissolved oxygen and cause biota kills<sup>34</sup>.

### DISCUSSION

*P. hoffmannianum* is similar in shape to *P. lima*, but larger and broader with dense areolae, and misidentified as *P. concavum* frequently, but can be distinguished by its ovoid shape and presence of areolae in the centre of the valve<sup>30</sup>. Both *P. hoffmannianum* and *P. belizeanum* have a prominent flared curved apical collar on the left valve bordering the periflagellar area<sup>35</sup>, and lack an apical spine of the periflagellar area<sup>17,35</sup>, although the latter species has a rounder and more prominent collar than the former one, and *P. hoffmannianum* has a more complex platelet configuration<sup>17,20,35</sup>, these two species were still considered as conspecific based on the subtle morphological overlaps found in cell shape, size, and ornamentation<sup>36</sup>. The areolae of *P. hoffmannianum* are distinct from the similar known species *P. ruetzlerianum*, *P. ruetzlerianum* has about 550 pentagonal-shaped areolae per valve while *P. hoffmannianum* has approximately 670 round to oval areolae<sup>17</sup>, although electron microscope was not performed in the present study, we still consider the current species (Fig. 2a) as *P. hoffmannianum*, for the cells of *P. ruetzlerianum* are round to ovoid with an average diameter of 28–35  $\mu\text{m}$ <sup>17</sup>.

*P. lima* can be distinguished by its size, shape, narrow periflagellar area and the presence of the valve and the marginal pores. The marginal pores can be used to differentiate *P. lima* at the light microscope level from the other species which are similar in shape, such as *P. concavum* or *P. compressum*<sup>37</sup>.

*P. micans* varies considerably in shape and size, can be confused with closely related species, such as *P. gracile*, *P. scutellum*, and *P. caribbaeum*. *P. gracile* has a strong winged apical spine, and the length-

width ratio usually larger than two, while *P. micans* usually smaller than two<sup>8</sup>; *P. scutellum* and *P. caribbaeum* are in the same size range as *P. micans*, but *P. scutellum* bears a shorter and broader apical spine, *P. caribbaeum* is heart-shaped and broadest around the anterior end, and *P. micans* is more tear-drop shaped and broadest around the middle<sup>15,35</sup>.

*P. rhathymum* was always considered as a synonym of *P. mexicanum*<sup>8</sup>, but some studies pointed out, the pores are arranged in organized pattern in *P. mexicanum* while disorganized in *P. rhathymum*<sup>38</sup>, and the former one has an apical wing-shaped spine terminating with two or three tips whereas the latter one only one simple spine<sup>10,11</sup>.

*P. sigmoides* was proposed as a synonym of *P. gracile*<sup>15</sup>, then treated as a distinct species for it has been consistently reported in Mexico<sup>9,13</sup>. *P. sigmoides* and *P. gracile* can be distinguished by the length-width ratio and the apical spine, for the former one, the length-width ratio usually larger than three, and the apical spine is on the right valve, sometimes sigmoid, while the latter one has a winged apical spine, sometimes like an arrow-head<sup>8,9</sup>, but it is still difficult to distinguish as either *P. sigmoides* or *P. gracile* for some cells, because they showed a sigmoid valve margin and a straight and long spine, or a slightly sigmoid spine with a straight valve margin, and mainly because there were no morphometric characters exclusive to *P. sigmoides*, it was suggested as a junior synonym of *P. gracile*<sup>13</sup>.

*P. hoffmannianum*, *P. lima*, and *P. rhathymum*, these three species occurred worldwide in the coastal areas, mostly in benthic and epiphytic habitats<sup>8</sup>, however, in this study, they were found in the water column, and although quantitative analysis of species populations was not performed, *P. rhathymum* showed a high concentration clearly. Actually, some literature indicate the benthic species can be pelagic, such as *P. lima*, this species has been found in substantial concentrations in the water column in Eastern Canada<sup>39</sup>, and high concentrations of this species were also found free-living in the subsurface waters of Northern Tunisia<sup>40</sup>, and on the other hand, we should note that, in the present study, all of the sites were chosen as shallow, about < 5 m deep areas, and these areas always suffering strong waves, and this may also explain why the benthic species appeared.

In this short report, five species of *Prorocentrum* were described from Sanya Bay, including three toxic species, *P. hoffmannianum*, *P. lima*, and *P. rhathymum*, and two red tide-forming species, *P. micans* and *P. sigmoides*. In particular, both

*P. hoffmannianum* and *P. lima* confirmed as the producers of okadaic acid and responsible for the DSP events<sup>19,25,40</sup>, while DSP is the most recurrent threat for the shellfish industries<sup>40</sup>, this information may indicate that, the coastal areas along Sanya Bay potentially face the danger of HABs, and close attention must be paid to the DSP events.

**Acknowledgements:** This work was supported by Hainan Provincial Natural Science Foundation of China (No. 417155), Scientific Research Foundation of Hainan Tropical Ocean University (No. RHDXB201618), and 2016 Institution of Higher Education Science Study Project in the Education Department of Hainan Province (No. Hnky2016ZD-4). We further acknowledge the valuable comments of Syuhei Ban and Shoko Tanabe.

## REFERENCES

- Hällfors H (2013) Studies on dinoflagellates in the northern Baltic Sea. PhD thesis, Univ of Helsinki.
- Simon N, Cras AL, Foulon E, Lemée R (2009) Diversity and evolution of marine phytoplankton. *CR Biol* **332**, 159–70.
- Hoppenrath M, Elbrächter M, Drebes G (2009) *Marine Phytoplankton. Selected Microphytoplankton Species from the North Sea around Helgoland and Sylt. E*, Schweizerbart Science Publishers, Stuttgart.
- Myat S, Thaw MSH, Matsuoka K, Lay KK, Koike K (2012) Phytoplankton surveys off the southern Myanmar coast of the Andaman Sea: an emphasis on dinoflagellates including potentially harmful species. *Fish Sci* **78**, 1091–106.
- Smayda TJ (2002) Adaptive ecology, growth strategies and the global bloom expansion of dinoflagellates. *J Oceanogr* **58**, 281–94.
- Dodge JD, Marshall HG (1994) Biogeographic analysis of the armored planktonic dinoflagellate *Ceratium* in the North Atlantic and adjacent seas. *J Phycol* **30**, 905–22.
- Gómez F (2005) A list of free-living dinoflagellate species in the world's oceans. *Acta Bot Croat* **64**, 129–212.
- Faust MA, Larsen J, Moestrup Ø (1999) Potentially toxic phytoplankton. 3. Genus *Prorocentrum* (Dinophyceae). In: Lindley JA (ed) *ICES Identification Leaflets for Plankton. Leaflet No 184*, International Council for the Exploration of the Sea, Denmark, pp 2–24.
- Hernández-Becerril DU, Cortes-Altamirano R, Alonso RR (2000) The dinoflagellate genus *Prorocentrum* along the coasts of the Mexican Pacific. *Hydrobiologia* **418**, 111–21.
- Gul S, Saifullah SM (2011) The dinoflagellate genus *Prorocentrum* (Prorocentrales, Prorocentraceae) from the North Arabian Sea. *Pak J Bot* **43**, 3061–5.
- Hoppenrath M, Chomérat N, Horiguchi T, Schweikert M, Nagahama Y, Murray S (2013) Taxonomy and phylogeny of the benthic *Prorocentrum* species (Dinophyceae)—A proposal and review. *Harmful Algae* **27**, 1–28.
- Steidinger KA, Tangen K (1997) Dinoflagellates. In: Tomas CR (ed) *Identifying Marine Phytoplankton*, Academic Press, London, pp 387–584.
- Cohen-Fernandez EJ, Castillo EMD, Ugarte IHS, Pedroche FF (2006) Contribution of external morphology in solving a species complex: The case of *Prorocentrum micans*, *Prorocentrum gracile* and *Prorocentrum sigmoides* (Dinoflagellata) from the Mexican Pacific coast. *Phycol Res* **54**, 330–40.
- Fritz L, Triemer RE (1985) A rapid simple technique utilizing calcofluor white M2R for the visualization of dinoflagellate thecal plates. *J Phycol* **21**, 662–4.
- Dodge JD (1975) The Prorocentrales (Dinophyceae). II. Revision of the taxonomy within the genus *Prorocentrum*. *Bot J Linn Soc* **71**, 103–25.
- Chihara M, Murano M (1997) *An Illustrated Guide to Marine Plankton in Japan*, Tokai Univ Press, Tokyo.
- Faust MA (1990) Morphologic details of six benthic species of *Prorocentrum* (Pyrrophyta) from a mangrove island, Twin Cays, Belize, including two new species. *J Phycol* **26**, 548–58.
- Morton SL, Faust MA (1997) Survey of toxic epiphytic dinoflagellates from the Belizean barrier reef ecosystem. *B Mar Sci* **61**, 899–906.
- Aikman KE, Tindall DR, Morton SL (1993) Physiology, potency of the dinoflagellate *Prorocentrum hoffmannianum* (Faust) during one complete growth cycle. In: Smayda TJ, Shimizu Y (eds) *Toxic Phytoplankton Blooms in the Sea*, Elsevier, Amsterdam, pp 463–8.
- Steidinger KA, Tangen K (1996) Dinoflagellates. In: Tomas CR (ed) *Identifying Marine Diatoms and Dinoflagellates*, Academic Press, New York, pp 387–584.
- Lu DD (2008) Phylum Dinzoa Cavalier-Smith, 1981. In: Liu JY (ed) *Checklist of Biota of Chinese Seas*, Science Press, Beijing, pp 175–86.
- Nagahama Y, Murray S, Tomaru A, Fukuyo Y (2011) Species boundaries in the toxic dinoflagellate *Prorocentrum lima* (Dinophyceae, Prorocentrales), based on morphological and phylogenetic characters. *J Phycol* **47**, 178–89.
- Faust MA (1991) Morphology of ciguatera-causing *Prorocentrum lima* (Pyrrophyta) from widely differing sites. *J Phycol* **27**, 642–8.
- Tindall DR, Miller DM, Bomber JW (1989) Culture and toxicity of dinoflagellates from ciguatera endemic regions of the world. *Toxicon* **27**, 83.
- Marr JC, Jackson AE, McLachlan JL (1992) Occurrence of *Prorocentrum lima*, a DSP toxin-producing species from the Atlantic coast of Canada. *J Appl*



- Phycol* **4**, 17–24.
26. Graneli E, Sundstrom B, Edler L, Anderson DM (1990) *Toxic Marine Phytoplankton*, Elsevier, New York.
  27. Uchida T (1977) Excretion of a diatom-inhibitory substance by *Prorocentrum micans* Ehrenberg. *Jap J Ecol* **27**, 1–4.
  28. Pinto JSD, Silva EDSE (1957) The toxicity of *Cardium edule* L. and its possible relation to the dinoflagellate *Prorocentrum micans* Ehr. *Jornal da Sociedade das Ciências Médicas de Lisboa* **121**, 304–26.
  29. Lassus P, Berthome JP (1988) *Status of 1987 Algal Blooms in IFREMER*, International Council for the Exploration of the Sea/Annex III C, Copenhagen.
  30. Fukuyo Y (1981) Taxonomical study on benthic dinoflagellates collected in coral reefs. *Bull Japan Soc Sci Fish* **47**, 967–78.
  31. Mohammad-Noor N, Daugbjerg N, Moestrup Ø, Anton A (2007) Marine epibenthic dinoflagellates from Malaysia - a study of live cultures and preserved samples based on light and scanning electron microscopy. *Nord J Bot* **24**, 629–90.
  32. Nakajima I, Oshima Y, Yasumoto T (1981) Toxicity of benthic dinoflagellates in Okinawa. *Bull Japan Soc Sci Fish* **47**, 1029–33.
  33. Lu S, Hodgkiss IJ (2004) Harmful algal bloom causative collected from Hong Kong waters. *Hydrobiologia* **512**, 231–8.
  34. Ajuzie CC, Houvenaghel GT (2009) Preliminary survey of potentially harmful dinoflagellates in Nigeria's coastal waters. *Fottea* **9**, 107–20.
  35. Faust MA (1993) *Prorocentrum belizeanum*, *Prorocentrum elegans* and *Prorocentrum caribbaeum*, three new benthic species (Dinophyceae) from a mangrove island Twin Cays, Belize. *J Phycol* **29**, 100–7.
  36. Herrera-Sepúlveda A, Medlin LK, Murugan G, Sierra-Beltrán AP, Cruz-Villacorta AA, Saavedra NYH (2015) Are *Prorocentrum hoffmannianum* and *Prorocentrum belizeanum* (DINOPHYCEAE, PROROCENTRALES), the same species? An integration of morphological and molecular data. *J Phycol* **51**, 173–88.
  37. Steidinger KA (1983) A re-evaluation of toxic dinoflagellate biology and ecology. *Prog Phycol Res* **2**, 147–88.
  38. Cortés-Altamirano R, Sierra-Beltrán AP (2003) Morphology and taxonomy of *Prorocentrum mexicanum* and reinstatement of *Prorocentrum rhathymum* (Dinophyceae). *J Phycol* **39**, 221–5.
  39. Bates SS (1997) Toxic phytoplankton on the Canadian east coast: implications for aquaculture. *Bull Aquacul Assoc Canada* **97**, 9–18.
  40. Sahraoui ID, Bouchouicha D, Mabrouk HH, Hlaili AS (2013) Driving factors of the potentially toxic and harmful species of *Prorocentrum* Ehrenberg in a semi-enclosed Mediterranean lagoon (Tunisia, SW Mediterranean). *Medit Mar Sci* **14**, 353–62.