

CAP-subgroups and $p\mathfrak{F}$ -hypercentrally embedded property

Yaxin Gao^{a,*}, Xianhua Li^b

^a Department of Mathematics, Changzhou University, Changzhou 213164 China

^b School of Mathematical Science, Guizhou Normal University, Guiyang 550025 China

*Corresponding author, e-mail: yxgaomath@163.com

Received 12 Mar 2024, Accepted 9 Sep 2025
Available online 11 Oct 2025

ABSTRACT: In this paper, we investigate $p\mathfrak{F}$ -hypercentrally embedded property of normal subgroups of a finite group and obtain some new results.

KEYWORDS: finite group, hypercentrally embedded, subgroup, Sylow subgroup

MSC2020: 20D10 20D20

INTRODUCTION

All groups in this paper are finite, G is always a finite group. π denotes a set of primes, G_π means a Hall π -subgroup of G , $O_\pi(G)$ is the largest normal π -subgroup of G , and $O^\pi(G)$ is the subgroup generated by all π' -elements of G . We use conventional notions and notations, as in [1, 2].

Recall that a class \mathfrak{F} of groups is called a formation if \mathfrak{F} is closed under taking homomorphic images and $G/(N_1 \cap N_2) \in \mathfrak{F}$ if $G/N_1, G/N_2 \in \mathfrak{F}$ for arbitrary normal subgroups N_1 and N_2 of G . A formation \mathfrak{F} is called saturated if $G/\Phi(G) \in \mathfrak{F}$ implies that $G \in \mathfrak{F}$. We use \mathfrak{U} to denote the formation of all supersolvable groups. It is clear that \mathfrak{U} is a saturated formation. A chief factor H/K of a group G is said to be \mathfrak{F} -central in G if $(H/K) \rtimes (G/C_G(H/K)) \in \mathfrak{F}$. A normal subgroup N of G is said to be $p\mathfrak{F}$ -hypercentrally embedded (resp., \mathfrak{F} -hypercentrally embedded) in G if either $N = 1$ or every p -chief factor of G (resp., chief factor of G) below N is \mathfrak{F} -central in G . The product of all normal $p\mathfrak{F}$ -hypercentrally embedded subgroups (resp., \mathfrak{F} -hypercentrally embedded subgroups) is called the $p\mathfrak{F}$ -hypercentre (resp., \mathfrak{F} -hypercentre) of G and denoted by $Z_{p\mathfrak{F}}(G)$ (resp., $Z_{\mathfrak{F}}(G)$). Clearly, a normal subgroup N of group G is $p\mathfrak{F}$ -hypercentrally embedded in G if and only if $N \leq Z_{p\mathfrak{F}}(G)$.

A subgroup H covers A/B if $HA = HB$ and avoids A/B if $H \cap A = H \cap B$, and has the cover-avoiding property in G if H either covers or avoids every chief factor of G (see [3]), in this case we may also say that H is a CAP-subgroup of G . A subgroup H of G is said to be semi cover-avoiding in G if there is a chief series $1 = G_0 < G_1 < \cdots < G_t = G$ of G such that for every $j = 1, 2, \dots, t$, H either covers G_j/G_{j-1} or avoids G_j/G_{j-1} (see [4]), in this case, H is also called a semi CAP-subgroup of G or partial CAP-subgroup of G in some literatures.

In recent years, many scholars have been interested in the influence of some property of the intersections between some subgroups and the subgroups

$O^p(G)$, or $O^p(G_p^*)$, or $G^{\mathfrak{F}}$ on the structure of a group G and give some criteria for p -supersolvability and p -nilpotency. For example, in [5], Guo and Isaacs investigated the supersolvability of a group G by assuming that $H \cap O^p(G) \triangleleft O^p(G)$ for any normal subgroup H of P with order d , where $P \in \text{Syl}_p(G)$ and d is a power of p with $1 \leq d < |P|$. In [6], the author proved that G is p -supersolvable if and only if $H \cap O^p(G_p^*)$ is s -permutable in G for all subgroups $H \leq P$ with $|H| = p^e$, where e is an integer with $e \geq 2$, $P \in \text{Syl}_p(G)$ and $|P| \geq p^{e+1}$. In addition, there are many literatures that have also investigated the influence of the properties of the intersections mentioned above on a group G (such as [7–11]), the authors obtained many results on p -supersolvability, p -nilpotency and p -supersolvable hypercenter of G . Our motivation is to develop such research by replacing the subgroups $O^p(G)$, or $O^p(G_p^*)$, or $G^{\mathfrak{F}}$ with general normal subgroups and obtain some new results. In [12], Lei, Li and Guo considered that the intersections between some subgroups and a normal subgroup satisfy permutability, their main theorems generalized many known results. In this paper, we continue to study this question and obtain some new characterizations for hypercentrally embedded property of normal subgroups of a finite group G by assuming that the intersections between some subgroups with fixed order and a normal subgroup are CAP-subgroups of G , which generalize the main theorem of [5].

PRELIMINARIES

In this section, for the sake of convenience, we present some basic results which will be used in the proofs of the Main Results section in this paper.

Lemma 1 ([13]) Let S be a CAP-subgroup of G and N a normal subgroup of G . Then

- (1) N is a CAP-subgroup of G ;
- (2) SN/N is a CAP-subgroup of G/N ;
- (3) SN is a CAP-subgroup of G ;

(4) $S \cap N$ is a CAP-subgroup of G .

Lemma 2 ([13]) Every minimal normal subgroup of G is a minimal CAP-subgroup of G .

Lemma 3 ([14]) Let G be a p -supersolvable group. Then the derived subgroup G' of G is p -nilpotent. In particular, if $O_{p'}(G) = 1$, then G has a unique (normal) Sylow p -subgroup.

Lemma 4 ([15]) Let C be a Thompson critical subgroup of a p -group P . Then the group $D := \Omega(C)$ is of exponent p if p is an odd prime, or exponent 4 if P is non-abelian 2-group. Moreover, every non-trivial p' -automorphism of P induces a non-trivial automorphism of D .

Lemma 5 ([16]) Let \mathfrak{F} be a solvably saturated formation and P a normal p -subgroup of G and C is a Thompson critical subgroup of P . If either $\Omega(C) \leq Z_{\mathfrak{F}}(G)$ or $P/\Phi(P) \leq Z_{\mathfrak{F}}(G/\Phi(P))$, then $P \leq Z_{\mathfrak{F}}(G)$.

Lemma 6 ([12]) Let \mathfrak{F} be a saturated formation, E be a normal subgroup of a finite group G and N a normal subgroup of G such that $N \leq \Phi(E)$. Then $E \leq Z_{p\mathfrak{F}}(G)$ if and only if $E/N \leq Z_{p\mathfrak{F}}(G/N)$.

MAIN RESULTS

In this section, we present the main results, which give some criterions for $p\mathfrak{F}$ -hypercentrally embedded property of subgroups.

Theorem 1 Let E be a normal subgroup of a finite group G and N a minimal normal subgroup of G contained in E and p be a prime divisor of $|G|$. Assume that $N_p > 1$ and there is a normal subgroup P of G_p such that $1 < N_p \leq P$ and $H \cap E$ is a CAP-subgroup of G for any subgroup H of P with order d , where $G_p \in \text{Syl}_p(G)$ and d is a power of p with $1 < d < |P|$. Then N is a p -group with $|N| \leq d$. Furthermore, if $|N| = d$, or $|N| < d$ and $N \not\leq \Phi(P)$, then $|N| = p$.

Proof: Since $N_p > 1$, we may let

$$1 < T_0 \leq T_1 \leq \dots \leq T_n = N_p \leq \dots \leq T_k = P \leq \dots \leq T_t = G_p$$

be a normal series of G_p such that $|T_i/T_{i-1}| = p$ for any $1 \leq i \leq t$. Assume that $|T_i| = d$. It is clear that $T_i \cap N = T_i \cap T_n \neq 1$. By the hypothesis and Lemma 1(4), $T_i \cap N = T_i \cap E \cap N$ is a CAP-subgroup of G . Then by Lemma 2 and the minimality of N , we have $T_i \cap N = N$. This implies that N is a p -group and $|N| \leq |T_i| = d$.

Assume $|N| = d$ and let $N \triangleleft U \leq P$. We can pick a subgroup U_1 of U such that $N \not\leq U_1$ and $|U_1| = d$. If not, then N is the unique maximal subgroup of U with order d and so $\Phi(U) = N$. This shows that U is cyclic. It follows that N is also cyclic and so $|N| = p$. By the hypothesis and Lemma 1(4), we get that $U_1 \cap N$ is a CAP-subgroup of G . By Lemma 2 and the minimality

of N , we have $U_1 \cap N = 1$ and so $|N| = |U_1| = p$, as required. Assume that $|N| < d$ and $N \not\leq \Phi(P)$. Then $n < i$ and $N \not\leq \Phi(T_{i+1})$ by [1, Hilfssatz III.3.3]. Hence we can pick a maximal subgroup K of T_{i+1} such that $N \not\leq K$. Clearly, $|K| = d$. It is easy to see that $K \cap N$ is a CAP-subgroup of G . Then by Lemma 2 and the minimality of N , we have $K \cap N = 1$ and so $|N| = p$. \square

Based on Theorem 1, we can obtain the following Theorem 2.

Theorem 2 Let \mathfrak{F} be a saturated formation containing \mathfrak{A} and E be a normal subgroup of a finite group G such that $G/E \in \mathfrak{F}$ and p be a prime divisor of $|G|$. Let K be a normal subgroup of G and $P \in \text{Syl}_p(K)$. Assume that $H \cap E$ is a CAP-subgroup of G for any normal subgroup H of P with order d , where d is a power of p with $1 \leq d < |P|$. Then either $K \leq Z_{p\mathfrak{F}}(G)$ or else $|P \cap E| > d$.

Proof: Let (G, E, K) be a counterexample with $|G| + |E| + |K|$ minimal order and let $T = P \cap E$. Then $K \not\leq Z_{p\mathfrak{F}}(G)$ and $|T| \leq d$. Assume that $O_{p'}(G) > 1$. By Lemma 1(2), it is easy to prove that $(G/O_{p'}(G), E/O_{p'}(G)/O_{p'}(G), KO_{p'}(G)/O_{p'}(G))$ satisfies the hypothesis of theorem. By the minimal choice of (G, E, K) , we have $KO_{p'}(G)/O_{p'}(G) \leq Z_{p\mathfrak{F}}(G/O_{p'}(G))$ or $|PO_{p'}(G)/O_{p'}(G) \cap EO_{p'}(G)/O_{p'}(G)| > d$, which implies that either $K \leq Z_{p\mathfrak{F}}(G)$ or $|P \cap E| > d$, a contradiction. Hence $O_{p'}(G) = 1$. Since $G/E \in \mathfrak{F}$, by [2, Proposition IV.1.5], every chief factor of G/E is \mathfrak{F} -central. In particular, every chief factor of G/E below KE/E is \mathfrak{F} -central. Let A/B be a chief factor of G such that $K \cap E \leq B \leq A \leq K$, then $B \cap E = A \cap E$ and so A/B is G -isomorphic to AE/BE . It is clear that $C_G(A/B) \leq C_G(AE/BE)$. Assume that $x \in C_G(AE/BE)$, then $[x, A] \leq [x, AE] \leq BE$. Since $A \leq G$, $[x, A] \leq A$. Hence $[x, A] \leq A \cap BE = B(A \cap E) = B$ and so $x \in C_G(A/B)$. Thus $C_G(A/B) = C_G(AE/BE)$. Since AE/BE is G -isomorphic to A/B ,

$$(A/B) \rtimes (G/C_G(A/B)) \cong (AE/BE) \rtimes (G/C_G(AE/BE)) \in \mathfrak{F}.$$

Hence $K/K \cap E \leq Z_{p\mathfrak{F}}(G/K \cap E)$. If $T = 1$, then $K \cap E$ is a p' -group. But $O_{p'}(G) = 1$, so $K \cap E = 1$. It follows that $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. This contradiction shows that $T \neq 1$. Let N be a minimal normal subgroup of G contained in T^G . Clearly, $N \leq T^G \leq E \cap K$. Since $|T| \leq d$ and $T \trianglelefteq P$, there is a normal subgroup H of P such that $T \leq H \leq P$ and $|H| = d$. Hence $T = P \cap E = H \cap E$ is a CAP-subgroup of G . If T avoids $N/1$, then $T \cap N = P \cap N = 1$. It implies that N is a p' -subgroup, which contradicts $O_{p'}(G) = 1$. Hence T must covers $N/1$, that is, $TN = T$, so $N \leq P \cap E$. This shows that $|N| \leq d$. Assume that $|N| = d$, then $N = P \cap E = T$ is a Sylow p -subgroup of $K \cap E$. It implies that $(K \cap E)/N$ is a p' -group. By the previous proof, we have $K/(K \cap E) \leq Z_{p\mathfrak{F}}(G/(K \cap E))$. Hence $K/N \leq Z_{p\mathfrak{F}}(G/N)$. Assume that $|N| < d$. By Lemma 1(2), we can get that $(G/N, E/N, K/N)$ satisfies the hypothesis. Hence $K/N \leq Z_{p\mathfrak{F}}(G/N)$ or $|P/N \cap E/N| > d/|N|$ by

the minimal choice of (G, E, K) . If $|P/N \cap E/N| > d/|N|$, then $|T| = |P \cap E| > d$, a contradiction. Hence $K/N \leq Z_{p\mathfrak{F}}(G/N)$ whether $|N| < d$ or $|N| = d$. If $N \leq \Phi(K)$, then by Lemma 6, we have $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. Hence we assume that $N \not\leq \Phi(K)$ and so $N \not\leq \Phi(P)$. Let M be any maximal subgroup of P . Since $M \cap E \trianglelefteq P$ and $|M \cap E| \leq |P \cap E| \leq d$, there is a normal subgroup H of P such that $M \cap E \leq H \leq M$ and $|H| = d$. Hence $M \cap E = H \cap E$ is a CAP-subgroup of G . By Theorem 1, $|N| = p$. Hence $N \leq Z_{p\mathfrak{F}}(G)$, which implies that $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. The contradiction completes the proof. \square

By Theorem 2, we can easily obtain the following corollaries.

Corollary 1 Let E be a normal subgroup of a finite group G such that G/E is supersolvable and p be a prime divisor of $|G|$ and let $P \in \text{Syl}_p(G)$. Assume that $H \cap E$ is a CAP-subgroup of G for any normal subgroup H of P with order d , where d is a power of p with $1 \leq d < |P|$. Then either G is p -supersolvable or else $|P \cap E| > d$.

Corollary 2 ([5]) Let p be a prime, let $P \in \text{Syl}_p(G)$, where G is a finite group, and let d be a power of p such that $1 \leq d < |P|$. Write $U = O^p(G)$, and assume that $H \cap U \trianglelefteq U$ for all subgroups $H \trianglelefteq P$ with $|H| = d$. Then either G is p -supersolvable or else $|P \cap U| > d$.

Remark 1 Corollary 2 represents one of the main theorems in [5]. It is easy to observe that this result can be directly derived from Theorem 2. Indeed, since $H \cap O^p(G) \trianglelefteq O^p(G)$ and $H \trianglelefteq P$, we have $H \cap O^p(G) \trianglelefteq G$. Clearly, $H \cap O^p(G)$ is a CAP-subgroup of G . Hence this result is a direct corollary of Theorem 2.

Theorem 3 Let \mathfrak{F} be a saturated formation containing \mathfrak{U} and E be a normal subgroup of a finite group G such that $G/E \in \mathfrak{F}$ and p be a prime divisor of $|G|$. Let K be a normal subgroup of G and $P \in \text{Syl}_p(K)$. Then $K \leq Z_{p\mathfrak{F}}(G)$ if one of the following holds:

- (1) $H \cap E$ is a CAP-subgroup of G for any subgroup H of P with order p . If P is a non-abelian 2-group, assume further that $H \cap E$ is a CAP-subgroup of G for any cyclic subgroup H of P with order 4.
- (2) $p = 2$, $|P| \geq 8$ and $H \cap E$ is a CAP-subgroup of G for any subgroup H of P with order 4.

Proof: Let (G, E, K) be a counterexample with $|G| + |E| + |K|$ minimal order. Similar to the proof of Theorem 2, we may assume that $O_{p'}(G) = 1$. By the hypothesis, $G/E \in \mathfrak{F}$ and $K \trianglelefteq G$. Similar to the proof of Theorem 2, it is easy to prove that $K/(K \cap E) \leq Z_{p\mathfrak{F}}(G/(K \cap E))$. If $P \cap E = 1$, then $K \cap E$ is a p' -group and so $K \cap E = 1$. It follows that $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. Hence we may assume that $P \cap E > 1$, so $K \cap E > 1$.

Firstly, we prove (1). Assume that $K \cap E < K$. Let K/T be a chief factor of G such that $K \cap E \leq T < K$. It is easy to see that (G, E, T) satisfies the hypothesis of the

theorem. Hence $K \cap E \leq T \leq Z_{p\mathfrak{F}}(G)$ by the minimal choice of (G, E, K) . Since $K/(K \cap E) \leq Z_{p\mathfrak{F}}(G/(K \cap E))$, $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. Hence $K \cap E = K$ and so $K \leq E$. Then every cyclic subgroup of P with order p and 4 (if P is a non-abelian 2-group) is a CAP-subgroup of G . Then by [13, Lemma 2.2(1) and (2)], every cyclic subgroup of P with order p and 4 (if P is a non-abelian 2-group) is a semi CAP-subgroup of K . By [17, Lemma 1.4], we can get that K is p -supersolvable. Since $O_{p'}(G) = 1$, by Lemma 3, we get $P \text{ char } K$ and so $P \leq G$. It is clear that (G, E, P) satisfies the hypothesis. If $P < K$, by the minimal choice of (G, E, K) , we have $P \leq Z_{p\mathfrak{F}}(G)$. Note that P is a Sylow p -subgroup of K , thus K/P is a p' -group. It follows that $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. This contradiction shows that $K = P$ is a p -group.

If K is a minimal normal subgroup of G , then $|K| = p$ and so $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. Hence K is not minimal normal in G . Now, let K/T be a chief factor of G . Then (G, E, T) satisfies the hypothesis of the theorem, and so $T \leq Z_{p\mathfrak{F}}(G)$ by the minimal choice of (G, E, K) . Since T is a p -group, $T \leq Z_{\mathfrak{F}}(G)$. Let L be any normal subgroup of G such that $L < K$. Then we also have $L \leq Z_{p\mathfrak{F}}(G)$. If $L \not\leq T$, then $K = TL \leq Z_{p\mathfrak{F}}(G)$, a contradiction. Hence we may assume that T is the unique normal subgroup of G such that K/T is a chief factor of G . If $|K/T| = p$, then $K/T \leq Z_{p\mathfrak{F}}(G/T)$ and so $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. Hence $|K/T| > p$. Let C be a Thompson critical subgroup of K . If $\Omega(C) < K$, we have $\Omega(C) \leq T \leq Z_{\mathfrak{F}}(G)$. Then, by Lemma 5, $K \leq Z_{\mathfrak{F}}(G) \leq Z_{p\mathfrak{F}}(G)$, a contradiction. This implies that $\Omega(C) = K$. Hence $\exp(K) = p$ or 4 by Lemma 4. Let x be an element of $K \setminus T$. Then $|\langle x \rangle| = p$ or 4.

By hypothesis, $\langle x \rangle \cap E$ covers or avoids K/T . In the former case, we have $\langle x \rangle K = \langle x \rangle T$, that is, $K = \langle x \rangle T$. It follows that K/T is cyclic and so $K/T \leq Z_{p\mathfrak{F}}(G/T)$, which implies that $K \leq Z_{p\mathfrak{F}}(G)$, a contradiction. In the latter case, $\langle x \rangle = \langle x \rangle \cap K = \langle x \rangle \cap T$ implies $\langle x \rangle = 1$ or $\Phi(\langle x \rangle)$, a contradiction. Thus, (1) holds.

Now, we prove (2). Let N be a minimal normal subgroup of G contained in $K \cap E$. By Theorem 1, N is a 2-group and $|N| \leq 4$. If $|N| = 4$, again by Theorem 1, $|N| = 2$, a contradiction. Hence $|N| = 2$ and so $N \leq Z(G)$. Let x be any element of $P \setminus N$ with order 2. Then $\langle x \rangle N = \langle x \rangle \rtimes N$ is a subgroup of P and $|\langle x \rangle N| = 4$. If $x \notin E$, then $\langle x \rangle \cap E = 1$ is a CAP-subgroup of G . If $x \in E$, then $\langle x \rangle N = \langle x \rangle N \cap E$ is a CAP-subgroup of G . Let A/B be any chief factor of G . Then $(\langle x \rangle N)A = (\langle x \rangle N)B$ or $\langle x \rangle N \cap A = \langle x \rangle N \cap B$.

Since N is a minimal normal subgroup of G , we distinguish three cases to complete the proof.

Case 1: $N \leq B \leq A$.

If $(\langle x \rangle N)A = (\langle x \rangle N)B$, then $\langle x \rangle A = \langle x \rangle B$; If $\langle x \rangle N \cap A = \langle x \rangle N \cap B$, then $(\langle x \rangle \cap A)N = (\langle x \rangle \cap B)N$. Note that $\langle x \rangle \cap N = 1$, hence $\langle x \rangle \cap A = \langle x \rangle \cap B$.

Case 2: $N \not\leq B$ and $N \leq A$.

Obviously, $N \times B = A$ and so $|A/B| = |N| = 2$.

If $(\langle x \rangle N)A = (\langle x \rangle N)B$, then we claim that $\langle x \rangle \cap A = \langle x \rangle \cap B$. If not, then $\langle x \rangle \cap B = 1$ and $\langle x \rangle \leq A$. It implies that $A = \langle x \rangle NB$ and so $|A/B| = 4$, a contradiction; If $\langle x \rangle N \cap A = \langle x \rangle N \cap B$, then $(\langle x \rangle \cap A)N = \langle x \rangle N \cap B$. If $\langle x \rangle \cap A \neq \langle x \rangle \cap B$, then $\langle x \rangle \cap B = 1$ and $\langle x \rangle \leq A$. Hence $\langle x \rangle N = \langle x \rangle N \cap B$ and so $\langle x \rangle N \leq B$. This shows that $\langle x \rangle \leq B$, a contradiction. Thus, $\langle x \rangle \cap A = \langle x \rangle \cap B$.

Case 3: $N \not\leq B$ and $N \not\leq A$.

If $(\langle x \rangle N)A = (\langle x \rangle N)B$, it is easy to prove that $\langle x \rangle \cap A = \langle x \rangle \cap B$. If not, we have $\langle x \rangle \cap B = 1$ and $\langle x \rangle \leq A$. Then $|AN| = |\langle x \rangle NB|$ and so $|A||N| = |\langle x \rangle||N||B|$. Hence $|A| = |\langle x \rangle||B|$. This induce that $\langle x \rangle A = A = \langle x \rangle B$; If $\langle x \rangle N \cap A = \langle x \rangle N \cap B$, we also get $\langle x \rangle \cap A = \langle x \rangle \cap B$. If not, $\langle x \rangle \cap B = 1$ and $\langle x \rangle \leq A$. Thus, $\langle x \rangle = \langle x \rangle (N \cap A) = \langle x \rangle N \cap B \leq B$, a contradiction.

The above three cases show that $\langle x \rangle \cap E = \langle x \rangle$ is a CAP-subgroup of G . Then by (1), $K \leq Z_{p\mathfrak{F}}(G)$. The proof is complete. \square

Based on the preceding theorems, we obtain the following Theorem.

Theorem 4 Let \mathfrak{F} be a saturated formation containing \mathfrak{U} , E be a normal subgroup of a finite group G such that $G/E \in \mathfrak{F}$ and p be a prime divisor of $|G|$. Let K be a normal subgroup of G , $P \in \text{Syl}_p(K)$ and d a power of p with $1 < d < |P|$. Assume that $H \cap E$ is a CAP-subgroup of G for any subgroup H of P with order d . If P is a non-abelian 2-group, assume further that $H \cap E$ is a CAP-subgroup of G for any cyclic subgroup H of P with order 4. Then $K \leq Z_{p\mathfrak{F}}(G)$.

Proof: By Theorem 3, we may assume that $d > p$. Similar to the proof of Theorem 2, we assume further that $O_{p'}(G) = 1$ and $K \cap E > 1$. Let N be a minimal normal subgroup of G contained in $K \cap E$. Clearly, $1 < N_p \leq P \trianglelefteq G_p$ for some Sylow p -subgroup G_p of G . By Theorem 1, N is p -group and $|N| \leq d$. If $|N| = d$, again by Theorem 1, $d = |N| = p$, a contradiction. Hence $|N| < d$.

Now, we claim that $K/N \leq Z_{p\mathfrak{F}}(G/N)$. Actually, if $d/|N| \neq 2$, then $(G/N, E/N, K/N)$ satisfies the hypothesis of the theorem, and so we can get $K/N \leq Z_{p\mathfrak{F}}(G/N)$ by induction. Assume $d/|N| = 2$. If P/N has a cyclic subgroup X/N of order 4 with $N \leq \Phi(X)$, then X is cyclic and therefore N is cyclic. Hence $|N| = 2$ and $d = 4$. By Theorem 3, $K \leq Z_{p\mathfrak{F}}(G)$. Hence, we may assume that $N \not\leq \Phi(X)$ for any cyclic subgroup X/N of order 4 in P/N . Then there is a maximal subgroup X_1 of X such that $X = X_1N$. Note that $|X_1| = d$, then $X/N \cap E/N = (X_1N \cap E)/N = (X_1 \cap E)N/N$ is a CAP-subgroup of G/N by Lemma 1 (2). This shows that K/N satisfies the hypothesis of Theorem 3, hence $K/N \leq Z_{p\mathfrak{F}}(G/N)$.

If $N \leq \Phi(K)$, then $K \leq Z_{p\mathfrak{F}}(G)$ by Lemma 6. If $N \not\leq \Phi(K)$, then $|N| = p$ by Theorem 1. Hence $K \leq Z_{p\mathfrak{F}}(G)$. \square

Corollary 3 Let G be a finite group, p be an odd prime divisor of $|G|$ and $P \in \text{Syl}_p(G)$ and let d be a power of p with $1 < d < |P|$. Assume that $H \cap O^p(G)$ is a CAP-subgroup of G for any subgroup H of P with order d . Then G is p -supersolvable.

CONCLUSION

In this paper, by investigating the intersections of some subgroups with fixed order and a normal subgroup satisfying the CAP-property, we obtain some new characterizations for the hypercentrally embedded property of normal subgroups of a finite group (i.e., Theorems 2–4). In particular, Theorem 2 generalizes one of the main theorems of [5].

Acknowledgements: The authors are grateful to the anonymous referees and editors for their valuable comments and specific corrections. This work was supported by the National Natural Science Foundation of China (Grant Nos. 12301019, 12261022 and 12401024).

REFERENCES

- Huppert B (1967) *Endliche Gruppen I*, Springer-Verlag, Berlin-Heidelberg-New York.
- Doerk K, Hawkes T (1992) *Finite Soluble Groups*, Walter de Gruyter, Berlin-New York.
- Guo X, Shum K (2003) Cover-avoidance properties and the structure of finite groups. *J Pure Appl Algebra* **181**, 297–308.
- Fan Y, Guo X, Shum K (2006) Remarks on two generalizations of normality of subgroups. *Chinese Ann Math Ser A* **27**, 169–176. [in Chinese]
- Guo Y, Isaacs I (2015) Conditions on p -subgroups implying p -nilpotence or p -supersolvability. *Arch Math* **105**, 215–222.
- Yu H (2017) Some sufficient and necessary conditions for p -supersolvability and p -nilpotence of a finite group. *J Algebra Appl* **16**, 1750052.
- Ballester-Bolinches A, Esteban-Romero R, Qiao S (2016) A note on a result of Guo and Isaacs about p -supersolvability of finite groups. *Arch Math* **106**, 501–506.
- Shen J, Qiao S (2018) p -supersoluble hypercenter and s -semipermutability of subgroups of a finite group. *Bull Iran Math Soc* **44**, 1185–1193.
- Miao L, Ballester-Bolinches A, Esteban-Romero R, Li Y (2017) On the supersoluble hypercentre of a finite group. *Monatsh Math* **184**, 641–648.
- Miao L, Li Y (2017) Some criteria for p -supersolvability of a finite group. *Commun Math Stat* **5**, 339–348.
- Guo X, Zhang B (2017) Conditions on p -subgroups implying p -supersolvability. *J Algebra Appl* **16**, 1750196.
- Lei D, Li X, Guo Y (2022) Weakly s -semipermutable subgroups and the $p\mathfrak{F}$ -hypercenter of finite groups. *Comm Algebra* **50**(11), 4610–4618.
- Li Y (2010) On cover-avoiding subgroups of Sylow subgroups of finite groups. *Rend Semin Mat Univ Padova* **123**, 249–258.
- Ballester-Bolinches A, Esteban-Romero R, Asaad M (2010) *Products of Finite Groups*, Walter de Gruyter, Berlin-New York.

15. Guo W, Skiba A (2015) Finite groups with generalized Ore supplement conditions for primary subgroups. *J Algebra* **432**, 205–227.
16. Chen X, Guo W (2016) On Π -supplemented subgroups of a finite group. *Comm Algebra* **44**, 731–745.
17. Ballester-Bolinches A, Esteban-Romero R, Li Y (2012) A question on partial CAP-subgroups of finite groups. *Sci China Math* **55**, 961–966.